

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6:

A61K 7/16

A1

(11) International Publication Number: WO 99/20237

(43) International Publication Date: 29 April 1999 (29.04.99)

(21) International Application Number: PCT/IB97/01634

(22) International Filing Date: 17 October 1997 (17.10.97)

(71) Applicant (for all designated States except US): ZAKRYTOE AKTSIONERNOE OBSCHESTVO OSTIM [RU/RU]; per. Sechenovsky, 6-3, Moscow, 119034 (RU).

(72) Inventors; and

(75) Inventors, and (75) Inventors, Applicants (for US only): RUDIN, Vsevolod Nikolaevich [RU/RU]; ul. Petrozavodskaya, 15-2-199, Moscow, 125502 (RU). ZUEV, Vladislav Petrovich [RU/RU]; ul. Tallinskaya, 2-316, Moscow, 123426 (RU). KOMAROV, Vladimir Fedorovich [RU/RU]; Shipilovsky proezd, 53-5-606, Moscow, 115582 (RU). MELIKHOV, Igor Vitallevich [RU/RU]; ul. Krupskol, 19-55, Moscow, 117331 (RU). MINAEV, Vladimir Vasillevich [RU/RU]; Naberezhnaya Tarasa Shevchenko, 5-471, Moscow, 121248 (RU). ORLOV, Andrei Yurlevich [RU/RU]; ul. Kantimirovskaya, 12-1-471, Moscow, 121357 (RU). MISHIN, Anatoly Aleksandrovich [RU/RU]; ul. Obrucheva, 3-2-77, Moscow, 117421 (RU). BOZHEVOLNOV, Viktor Evgenievich [RU/RU]; ul. Bolshaya Kommunisticheskaya, 24/28-45, Moscow, 109004 (RU).

(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

Published

With international search report.

(54) Title: STOMATIC COMPOSITION

(57) Abstract

This invention relates to the field of medicine, and in particular to the stomatology and may be used for preventive treatment and combatting caries, parodenitis and paradentosis. The stomatic composition on the basis of hydroxyapatite also comprises abrasive materials, humectants, thickeners, surfactants, flavouring agents, and a number of optional ingredients. The composition comprises hydroxyapatite in the form of particles of ultra finely divided hydroxyapatite with an average size by length (1), width (d) and thickness (h) of about 1 x d x h = 0.06 μ m x 0.015 μ m; x 0.005 μ m is capable to stimulate reparative osteogenesis processes and possessing a high bioactivity and specific pharmacological activity. The ultra finely divided hydroxyapatite is present in the composition in the amount of 0.1 % to 50 % by weight. The ultra finely divided hydroxyapatite hydroxyapatite which contains about 99.9 % of Ca₁₀(PO₄)₆(OH)₂ by weight. The stomatic composition can be used to cure microdefects of the basic substance of the dental enamel, can be applied to prepare toothpastes, toothcreams and gels and can also be included as a component of chewing gum, pastilles, tooth elixir, formulations to rinse mouth.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia .	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
ΑU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
ΑZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of Americ
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	Li	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

WO 99/20237 PCT/IB97/01634

STOMATIC COMPOSITION

Description

Field of the Invention

This invention relates to the field of medicine, and in particular to the field of stomatology and may be used for preventive treatment and curing of caries, parodenitis and paradentosis.

Prior art

For these above-captioned purposes, stomatic compositions comprising hydroxyapatite (HA) have found an extensive application in the stomatologic practice.

There are certain compositions having a favourable effect including synthetic HA containing 92 to 97% $Ca_{10}(PO_4)_6(OH)_2$, 3 to 6% H_2O and 0,3% $CaCO_3$ with an average particle size of 1 to 15 μm .

Such a stomatic composition, for instance, according to Patent EP 0344832 cl. A61K 7/16, comprises save the stated HA, water-soluble casein material or sodium trimetaphosphate, as an anti-caries agent and also other well known ingredients which depend upon the forms of the product manufactured, such as various humectants, binding thickeners, surfactants, flavouring agents.

The known stomatic composition (EP 03442746 cl. A61K 7/18, publ.23.11.89) supplementary includes a fluorine-containing compound in the form of NaP or sodium monophluorphosphate as an anti-caries agent.

The amount of HA present in the stomatic composition is in the range of 1 to 50%, usually 2 to 20% by weight of the stomatic composition. The stomatic composition comprises some other ingredients: humectants, thickeners, surfactants

and flavouring agents commonly known to those skilled in the art in all formulations of such products.

However, the stomatic compositions stated possess a relatively poor anti-caries effect and is not useful in the preventive medicine and in the treatment of inflammatory-destructive diseases of paradentium tissues.

Disclosure of the invention

It is an object of the present invention to create a stomatic composition comprising compounds capable to cure microdefects of the basic substance of the dental enamel to combat caries developing (e.g. to provide an anti-caries activity) and to prevent the spread of such inflammable-destructive diseases of paradentium tissues as paradenitis and paradentosis, and also compounds capable to stimulate reparative osteogenesis processes and possessing high bioactivity and specific pharmacological activity.

It is a further object of the invention to create a stomatic composition being identic to the basic substance of the dental enamel in its substance contents and crystalline parameters, as the acid formed in the materials covering dental surfaces causes destruction of mineral hydroxyapatyte out of which enamel is composed and has a result due to which calcium ion loss occurs.

The task surprisingly has been solved in a composition as defined in claim 1.

A preferred composition having a more pronounced effect in view of the improvements obtained according to the invention comprises particles of hydroxyapatite with an average particle size in length (l), width(d) and thickness(h) of about $l=0.06~\mu m$ +/- 50 %, $d=0.015~\mu m$ +/- 50 % and $h=0.005~\mu m$ +/- 50 %.

A most preferred composition having a surprisingly superior effect in view of the improvements obtained according to the invention comprises particles of hydroxyapatite with an average particle size in length (1),

width(d) and thickness(h) of 1 about 0,06 μ m, d about 0,015 μ m, h about 0,005 μ m.

Being introduced into the composition, HA possesses osteo-reparative properties and contains preferably about 100% $Ca_{10}(PO_4)_6(OH)_2$.

The specific surface of HA used in the composite advantageously is $100 \text{ to } 150 \text{ m}^2/\text{g}$.

The amount of HA present in the oral composition of the present invention is in the range of 0,1% to 50%, preferably from 0,1% to 25%, and most preferably from about 0,2% to 20% by weight of the oral composition.

The composition reacts to a change in the biochemical environment, for instance a rapid dissolvement of ultra finely divided HA occurs when the pH is decreasing, that provides an active utilization of Ca and PO_4 - ions in the osteogenesis process: the size and configuration of the inventive crystals are adapted to the maximum to the dental enamel structure, which is mostly composed of HA, that suggests its use in the osteo-reparative process as a building material.

The ultra finely divided HA possesses a high adhesive-sorption activity to the dental enamel and to microdefects on its surface, that favour the preventive measures preventing the spread of caries disease and also possesses a high sorption activity in respect to proteins and aminoacids of paradentium tissues, that stimulates an active preventive treatment of the inflammable-destructive diseases such as paradenitis and paradentosis.

Moreover, the stomatic composition of the present invention will contain other conventional ingredients in addition to HA possessing osteo-reparative properties, whose introduction into the composition depends on the form of the product. For instance, in the case of an oral product in the form of dentifrice paste, cream or gel, the product will comprise a liquid phase containing humectants and binding thickeners which act to maintain the particulate solid

abrasive and HA crystals in the form of stable suspension in the liquid phase.

Surfactants and flavouring agents are also usual ingredients for various inventive embodiments of oral compositions.

The humectants usually used are glycerol or sorbitol. However, other humectants may be used according to the invention including polyethyleneglycol, propyleneglycol, lactitol and hydrogenated corn syrop. The amount of humctant will generally range from about 0% to 85% by weight of product. The remainder of the liquid phase will consist substantially of water. The liquid phase can be water or a non-aqueous composition.

As binding agents and thickeners, various substances can be used such as sodium carboxymethylcellulose, sodium hydroxyethylcellulose and xanthan gum. Natural gum bindings can be included such as gum tragacanth, gum karaya of Irish moss, etc. Any mixture of binding agents and thickeners can be also used. The amount of bindings and thickeners usually included into the oral composition is in the range of 0% to 10% by weight of the oral composition.

Moreover, any materials as widely disclosed in the literature generally also might be used for the invention as surfactants, i.e. surfactants like sodium lauryl sulphate, dodecylbenzene sulphonate and sodium lauryl sarcosinate. Other anionic surfactants also can be used as well as cationic and amphoteric and non-ionic surfactants. Surfactants are generally present in the composition in the amount of 0% to 5% by weight of the oral composition.

Flavours that are generally used in the oral compositions are those based on oils of spearmint and peppermint and might be used for the invention. Examples of other flavouring materials used are menthol, clove, wintergreen, eucalyptus and aniseed. A preferable amount of flavours is from 0% to 5% by weight in respect to the oral composition.

As abrasive materials, silica dioxide of various modifications, aluminium oxide, calcium carbonate, dicalcium phosphate anhydrite, dicalcium phosphate dihydrate, sodium metaphosphate insoluble in water, and thereof mixtures may be used. The amount of abrasive materials ranges from 0.0% to 25%. The oral composition may include a wide variety of optional ingredients. These include antimicrobial and antiplaque agents for example chlorhexidine or 2,4,4-trichloro-2hydroxy-diphenyl ether, or zink compounds (see EPA-161898) anti-tartar ingredients such as condensed phosphates, e.g. alkali et al pyrophosphates, hexametaphosphatesor polyphosphates (see US-A-4 515772 and US-A-4 627977) or zink citrates (see US-A-4 100269), sweetening agents such as saccharin. Preservatives such as formalin, sodium benzoate. colouring agents (for instance titanium dioxide) or pHcontrolling agents, such as acid base or buffer agents the oral composition may also include agents enhancing the gingivitis system of the mouth cavity and representing extracts of various natural plants such as urtica, millefolium, chamomilla hypericum, salvia, etc. in the aqueous or aqueous-alcoholic forms.

The stomatic composition depending on its form (dentifrice paste, cream or gel) is maintained in contact with the tissue of the oral cavity from 15 sec to 12 hours.

The following examples of dentifrice pastes and gel comprising synthetic ultra finely divided HA possessing osteo-reparative properties as described above illustrate the invention. Percentages and parts of the components are by weight.

Belowstanding preferred embodiments of the invention are shown in its composition.

Examples N1 and 2.

Toothpaste prepared from the following ingredients.

Ingredi	0,2 2,0 22,0 15,0 ylcellulose 1,0 1,0 20,0 20,0 20,0 17,0				
Example	1	2			
Ultra finely divided					
Hydroxyapatite	0,2	2,0			
Silica aerogel	22,0	15,0			
Sodium carboxymethylcellulose	1,0	1,0			
Glycerol distilled	20,0	20,0			
Sorbitol	20,0	17,0			
Titanium dioxide	0,6	0,5			
Sodium benzoate	0,4	0,6			
Aqueous-alcohol extract					
of chamomilla	1,0	0,8			
Aqueous-alcohol extract					
of hypericum	1,0	0,8			
Sodium saccharin	0,1	0,06			
Flavour	1,0	1,3			
Sodium lauryl sulphate	Sodium lauryl sulphate 1,5 1,5				
Water	to 100,0	to 100,0			

Examples N 3 to 7 Toothpaste prepared from the following ingredients.

Ingredients,%						
Example	3	4	5	6	7	
Ultra finely divided						
hydroxyapatite	2,5	2,5	2,5	2.5	2.5	
Silica aerogel			17,0			
Sodium					·	
hydroxyethylcellulose	1,6	-	-	1,6	-	
Sodium						
carboxymethylcellulose	-	1,1	1,1	_	0,9	
Sorbitol	20,0	20,0	16,0	20,0	20,0	
Glycerol distilled	20,0	20,0	20,0	20,0	20,0	
Polyethyleneglycol	-	-	5,0	-	-	
Sodium lauryl sulphate	1,5	1,5	1,5	1,5	1,5	
Tetrasodium pyrophosphat	e -	1,5		-	-	
Tetrapotassium						
pyrophosphate	-	-	-	2,5	-	
Sodium trimetaphosphate	-	-	2,0		-	
Zinc citrate trihydrate	-	-	-	-	0,5	
Titanium dioxide	0,6	0,6	0,6	0,6	0,6	
Sodium benzoate	0,5	0,5	0,6	-	-	
Formalin	-	-	-	0,05	0,05	
Aqueous-alcohol extract						
of salvia	0,5	0,5	-	-	-	
Aqueous-alcohol extract						
of millefolium	0,9	0,9	0,5	0,5	-	
Aqueous-alcohol extract			٠			
of chamomilla	-	-	1,0	0,7	-	
Triclosan	_	-	0,2	-	0,2	
Sodium saccharin	0,06	0,06	0,06	0,06	0,06	

R

Flavour	1,0	1,0	1,0	1,0	1,0
Water		in	all examp	e to	100,0

Examples N8 and 9
Gel preventing paradenitis.

Ing		
Example	8	9
Ultra finely divided		
hydroxyapatite	5,0	4,0
Sodium hydroxyethylcellulose	2,0	2,5
Silica aero	5,0	-
Glycerol distilled	10,0	-
Sorbitol	25,0	45,0
Sodium benzoate	0,5	-
Triclosan	-	0,3
Flavour	0,2	0,15
Sodium lauryl sulphate	0,2	0,15
Sodium saccharin	0,07	0,07
Water	to 100,00	to 100,00

Industrial application

The stomatic composition can be used to cure microdefects of the basic substance of the dental enamel, e.g. to prevent the spread of caries, and is also useful for

preventive measures avoiding the spread of inflammabledestructive diseases of paradentium tissues, such as pardenitis and paradentosis.

The stomatic composition can be used in the form of tooth pastes, tooth creams and gels. Moreover, the composition can be included as a component in jewing gum, pastilles, tooth elixir and formulations to rinse mouth.

The stomatic composition according to the invention is capable to stimulate reparative osteogenesis processes and possessing a high bioactivity and specific pharmacological activity. Moreover, this composition is useful for combatting dental caries and to prevent the spread of such inflammable-destructive diseases of paradentium tissues as paradenitis and paradentosis, on the basis of hydroxyapatite also optionally comprising abrasive materials, humectants, thickeners, surfactants, flavouring agents, and a number of optional ingredients.

Claims:

- 1. A stomatic composition characterised in that it comprises particles of hydroxyapatite with an average particle size in length (1), width (d) and thickness (h) of: 1 from about 0,2 μm to about 0,01 μm , d from about 0,1 μm to about 0.001, and h from about 0.1 μm to about 0,001 μm .
- 2. The stomatic composition according to claim 1 characterised in that it comprises particles of hydroxyapatite with an average particle size in length (l), width(d) and thickness(h) of about l = 0,06 μ m +/- 50 %, d = 0,015 μ m +/- 50 % and h = 0,005 μ m +/- 50 %.
- 3. The stomatic composition according to claim 1 characterised in that it comprises particles of hydroxyapatite with an average particle size in length (1), width(d) and thickness(h) of about 1 = 0,06 μ m, d = 0,015 μ m, h = 0,005 μ m.
- 4. A stomatic composition characterised in that it comprises particles of hydroxyapatite having a specific surface of hydroxiapatite from about 100 m 2 /g to about 150 m 2 /g.
- 5. The stomatic composition according to one of claims 1 to 4 characterized in that it comprises said hydroxyapatite particles ultra finely divided.

- 6. The composition according to one of claims 1 to 5 characterised in that the ultra finely divided hydroxyapatite particles are present in the composition in an amount of 0,1% to 50% by weight.
- 7. The composition according to one of claims 1 to 6 characterised in that the ultra finely, divided hydroxyapatite is a synthetic hydroxyapatite which contains 99,9% of $Ca_{10}(PO_4)_6(OH)_2$ by weight.
- 8. The composition according to one of claims 1 to 7 further characterised by at least one substance of the group consisting of
 - humectants in a range from about 0% to 85% by weight,
 - bindings and thickeners a range of 0% to 10% by weight,
 - abrasive materials in a range from 0.0% to 25%,
 - Surfactants in a range from 0% to 5% by weight,
 - Flavours in a range from 0% to 5% by weight.
- 9. The composition according to one of claims 1 to 8 further characterised by agents enhancing the gingivitis system of the mouth cavity and comprising extracts of natural plants including at least one of the group consisting of urtica, millefolium, chamomilla hypericum, salvia, etc. in the aqueous an in the aqueous-alcoholic form.
- 10. The composition according to one of claims 1 to 9 further characterised by antimicrobial and anti-plague agents.

 $\frac{N_{q_1}}{N}$

INTERNATIONAL SEARCH REPORT

national Application No

		i i				
A. CLASSI IPC 6	FICATION OF SUBJECT MATTER A61K7/16					
According to	o International Patent Classification(IPC) or to both national classifica	ation and IPC				
	SEARCHED					
Minimum do IPC 6	ocumentation searched (classification system followed by classification $A61K$	on symbols)				
Documental	tion searched other than minimum documentation to the extent that so	uch documents are include	d in the fields searched			
Electronic d	ata base consulted during the international search (name of data bas	se and, where practical, se	arch terms used)			
C. DOCUM	ENTS CONSIDERED TO BE RELEVANT					
Category 3	Citation of document, with indication, where appropriate, of the rele	evant passages	Relevant to claim No.			
X	EP 0 786 245 A (SANGI CO., LTD) 3 1997 see the whole document	1-10				
X	EP 0 664 133 A (AKTIONERNOE OBSCH ZAKRYTOGO TIPA "OSTIM") 26 July 1 see claim 1; example 1	1				
E	WO 98 18719 A (AKTSIONERNOE OBSCH ZAKRYTOGO TIPA "OSTIM") 7 May 199 see claims 1,14 	1				
Furth	ner documents are listed in the continuation of box C.	X Patent family me	mbers are listed in annex.			
° Special cal	tegories of cited documents :		ned after the international filing date			
"A" document defining the general state of the art which is not considered to be of particular relevance or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention						
filling date cannot be considered novel or cannot be considered to						
which i citation	"L" document which may throw doubts on priority claim(s) or which is cited to establish the publicationdate of another citation or other special reason (as specified) "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the					
other n	"O" document referring to an oral disclosure, use, exhibition or other means document is combined with one or more other such document other means, such combination being obvious to a person skilled in the art.					
	an the priority date claimed actual completion of theinternational search	"&" document member of Date of mailing of the	international search report			
9	July 1998	16/07/19	98			
Name and m	nailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2	Authorized officer				
٠	NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016	Glikman,	J-F			

INTERNATIONAL SEARCH REPORT



PCT/IB 97/01634

	Patent document cited in search report		Publication date		atent family nember(s)	Publication date	
EP	786245	Α	30-07-1997	JP	9202717 A	05-08-1997	
EP	664133	Α	26-07-1995	RU WO	2077329 C 9503074 A	20-04-1997 02-02-1995	
WO	9818719	Α	07-05-1998	NONE			